


# PRV

PATENT- OCH REGISTRERINGSVERKET  
Patentavdelningen

PC1/6M 2003 / 001385 107389654



## Intyg Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

Ansökan ingavs ursprungligen på engelska.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

The application was originally filed in English.

(71) Sökande AstraZeneca AB, Södertälje SE  
Applicant (s)

(21) Patentansökningsnummer 0201005-6  
Patent application number

(86) Ingivningsdatum 2002-04-02  
Date of filing

Stockholm, 2003-02-19

För Patent- och registreringsverket  
For the Patent- and Registration Office

Kerstin Gerdén

Avgift  
Fee 170:-

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

## BEST AVAILABLE COPY

PATENT- OCH  
REGISTRERINGSVERKET  
SWEDEN

Postadress/Adress  
Box 5055  
S-102 42 STOCKHOLM


Telefon/Phone  
+46 8 782 25 00  
Vx 08-782 25 00

Telex  
17978  
PATOREG S

Telefax  
+46 8 666 02 86  
08-666 02 86

## NEW PROCESS

The present invention relates to an improved process for the preparation of the compound 2-ethoxy-3-[4-(2-(4-methanesulfonyloxyphenyl)ethoxy)phenyl]propanoic acid, as shown  
5 in formula I below



10 or the (R) or the (S) enantiomer thereof, or a pharmaceutically-acceptable salt thereof, and solvates thereof.

The above compound is intended for therapeutic use in the Insulin Resistance Syndrome (IRS) including type 2 diabetes mellitus, which refers to a cluster of manifestations  
15 including insulin resistance with accompanying hyperinsulinaemia, possible type 2 diabetes mellitus, arterial hypertension, central (visceral) obesity, dyslipidaemia observed as deranged lipoprotein levels typically characterised by elevated VLDL (very low density lipoproteins), small dense LDL particles and reduced HDL (high density lipoprotein) concentrations and reduced fibrinolysis.

20

Recent epidemiological research has documented that individuals with insulin resistance run a greatly increased risk of cardiovascular morbidity and mortality, notably suffering from myocardial infarction and stroke. In type 2 diabetes mellitus atherosclerosis related conditions cause up to 80% of all deaths.

25


In clinical medicine there is awareness of the need to increase the insulin sensitivity in IRS suffering patients and thus to correct the dyslipidaemia which is considered to cause the

accelerated progress of atherosclerosis. However, currently this is not a universally well defined disease.


The compound of formula I is disclosed in PCT Publication Number WO99/62872. Two  
5 alternative processes are disclosed for the preparation of the compound of formula I in  
Examples 1 and 2 of the application. We have discovered an improvement in relation to  
one of the processes disclosed.

Specifically we have found an improved method for the synthesis of the penultimate  
10 esterified intermediate, the final step of the reaction being the conversion of the ester group  
into the acid of the final product.


Accordingly the present invention provides a process for the preparation of a compound of formula I



in which R represents H or an acid protecting group which comprises reacting a compound of formula II



in which R is as previously defined with a compound of formula III



wherein X is a suitable leaving group in the presence of a base using water as the diluent.

- 5 Suitably the process is carried out at a temperature in the range 0°C to 250°C and preferably in the range 50°C to 150°C. More preferably the process is carried out at a temperature in the range of 80°C to 130°C and most preferably in the range of 90°C to 110°C.

The term "acid protecting group" means that the acid is protected from reaction by  
10 forming a suitable acid derivative such as an ester or amide or by other means of protection of carboxylic acid groups known in the art. Examples of suitable means of protection and acid derivatives (as well as means of formation and eventual deprotection), may be found in T.W. Greene and P.G.M. Wuts, "Protective Groups in Organic Synthesis", Third Edition, John Wiley & Sons, New York, 1999. The nature of the ester is not important in  
15 the performance of the process since its function is to act as a protecting group. The improvements relate to the application of phase transfer catalysis to the process. Preferably R is H, benzyl or a (1-4C)alkyl group, such as methyl, ethyl or propyl. More preferably R is a (1-4C)alkyl group. Most preferably R is ethyl.

- 20 Suitably X is halo, for example bromo, chloro or iodo, an optionally substituted phenylsulfonyloxy group, in particular (4-methylphenyl)sulfonyloxy group or 2,4,6-triisopropylphenylsulfonyloxy group or an alkylsulphonyloxy group for example methanesulphonyloxy. Preferably X is a methanesulphonyloxy group.

- 25 Suitable bases include carbonates, hydrogen carbonates or hydroxides particularly of alkali metals. Preferably the base is sodium carbonate, sodium hydrogen carbonate, potassium carbonate or potassium hydrogen carbonate.


Suitably the molar ratio of the compound of formula III to the compound of formula II is in the range of 0.5 to 10, preferably in the range 0.8 to 4 and more preferably in the range of 1.0 to 3 and most preferably is in the range of 1.2 to 1.6.

Suitably the molar ratio of the base to the compound of formula II is in the range of 0.5 to 10, preferably in the range 0.8 to 4 and more preferably in the range of 1.0 to 3 and most preferably is in the range of 1.2 to 1.6.

The process of the invention has the following advantages. The reaction times are more rapid than the reactions known in the prior art and therefore the process is less costly to run. In addition the process gives higher yields and the product is of a higher purity than previously disclosed processes for the preparation of the compound of formula I. Also, it is a considerable advantage in terms of waste disposal, environmental reasons and cost to use water as the diluent instead of an organic liquid. Further, the process is consistently reproducible and robust.


Converting the acid ester derivative may be accomplished simply by hydrolysis (acidic or alkaline or enzymatic) of the ester to the acid, such a step being known to the skilled person, such as described in the examples below and in Example 2 i) of WO99/62872.

In another aspect the present invention provides a process for the preparation of a compound of formula I




in which R represents H which comprises reacting a compound of formula II

5



in which R represents an acid protecting group with a compound of formula III



- 5 wherein X is a suitable leaving group in the presence of a base and using water as the diluent to give a compound of formula I in which R is an acid protecting group and then removing the protecting group to give a compound of formula I in which R is H.

In a preferred aspect the protecting group is an ester and the protecting group removal step  
 10 comprises a hydrolysis step. The hydrolysis step may be acid or base catalysed. Optionally an organic liquid may be present in the hydrolysis step for example acetone, 2-butanone, methanol, ethanol, tetrahydrofuran or dioxane.

In a preferred aspect the process provides the S-enantiomer of the compound of formula I in  
 15 which R is H by using the S-enantiomer of the compound of formula II in which R represents H or an acid protecting group followed by hydrolysis when R is an acid protecting group.

The compound of formula I in which R is H may be purified by recrystallisation. Suitable  
 20 recrystallisation solvents include one or more of the following ethanol, water, isopropyl acetate, isopropanol, isooctane and toluene.

The invention is illustrated by the following non-limiting examples.

### Abbreviations

5 EtOAc = ethyl acetate

HPLC = high-pressure liquid chromatography

i-PrOAc = isopropyl acetate

EtOH= ethanol

### 10 Preparation of Starting Material

2-(4-Methanesulfonyloxyphenyl)ethylmethanesulfonate was prepared as described in WO99/62872.

### 15 Example 1

#### (S)-2-ethoxy-3-[4-(2-[4-methanesulfonyloxyphenyl]ethoxy)phenyl] propanoic acid

A mixture of 2-(4-methanesulfonyloxyphenyl)ethylmethanesulfonate (9.88 g / 33.57 mmol), ethyl (S)-2-ethoxy-3-(4-hydroxyphenyl)propanoate (5.0 g / 20.98 mmol), Na<sub>2</sub>CO<sub>3</sub> (2.96 g / 27.9 mmol eq) and water (10ml) was boiled under reflux with vigorous stirring for 4.5 h. The mixture was cooled to 2°C and acetone (13.5ml) was added. The mixture containing ethyl (S)-2-ethoxy-3-[4-(2-[4-methanesulfonyloxy-phenyl]ethoxy)phenyl] propanoate was allowed to warm up to ambient temperature.


25 The acetone solution was kept at T<sub>i</sub> = 25 °C in the reactor. LiOH·H<sub>2</sub>O powder (1.144 g / 27.27mmol) dissolved in water (20ml) was charged continuously over 30 minutes under vigorous stirring at 25±5 °C. The reaction was continued for 7 hours. The reaction was quenched by addition of EtOAc (1.5 ml) at T<sub>i</sub> = 25±5 °C.

The identity of the product was confirmed by HPLC and gave a relative area of 76%.


30

**Claims**


1. A process for the preparation of a compound of formula I



5 in which R represents H or an acid protecting group which comprises reacting a compound of formula II




10 in which R is as previously defined with a compound of formula III




wherein X is a suitable leaving group in the presence of a base and using water as a diluent.

15 2. A process for the preparation of a compound of formula I






in which R represents H which comprises reacting a compound of formula II



in which R represents an acid protecting group with a compound of formula III



wherein X is a suitable leaving group in the presence of a base and using water as the diluent to give a compound of formula I in which R is an acid protecting group and then removing the protecting group to give a compound of formula I in which R is H.

3. A process according to claim 2 in which the acid protecting group is removed by hydrolysis.

4. A process according to any preceding claim in which R is H, benzyl or a (1-4C)alkyl group.

5. A process according to any preceding claim in which X is halo, an optionally substituted phenylsulfonyloxy group or an alkylsulphonyloxy group.

6. A process according to any preceding claim in which the base is selected from carbonates, hydrogen carbonates or hydroxides of alkali metals.


7. A process according to any preceding claim in which the compound of formula I is the S enantiomer.

100679-1 SE

**Abstract**


A process for the preparation of a compound of formula I

suitable the




5

in which R represents H or an acid protecting group which comprises reacting a compound of formula II



10

in which R is as previously defined with a compound of formula III



wherein X is a suitable leaving group in the presence of a base and using water as a diluent.

15

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**